

Disclaimer

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.
- Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

Hyalein[®]S Mini ophthalmic solution

Hyalein S Mini ophthalmic solution can be used while wearing any type of contact lens except colored contact lenses.

Soft

O₂

Hard

Disposable

This product can also be used by people not wearing contact lenses.

* Because some eye drops cannot be used while wearing soft contact lenses, read the instructions on the package or package insert carefully.



Precautions for use



When not to use the product

(Not observing the following precautions may aggravate present symptoms or cause adverse reactions.)

Do not use the product if you are one of the following people.

- (1) Patients who have experienced allergic symptoms associated with drugs, etc
- (2) People who have received the following diagnoses:
Dry eye, Sjögren syndrome, Stevens-Johnson syndrome, or corneal infection
- (3) People who have the following symptoms:
Sudden visual deterioration or severe pain in the eye(s)
(Do not self-administer any treatment on your own judgment, as doing so may cause symptoms to worsen. Please consult a doctor.)



Consultation

1. The following people should consult with their doctor, pharmacist, or registered pharmaceutical distributor **before using this product**.
 - (1) People receiving treatment from a doctor
 - (2) People who have experienced an allergic reaction to medicine or other substances
 - (3) People who have the following symptoms in addition to symptoms of the eyes:
 - Dryness of the mouth or dryness of the nasal cavity
 - High fever, lip erosion, sore throat, persistence of or sudden worsening of widespread rash/redness of the skin
 - (4) People who have received the following diagnosis: Glaucoma
2. If you experience any of the following symptoms **after using this product**, it may be an adverse reaction. In such a case, please stop using the product immediately, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.

Area affected	Symptom
Skin	Rash/redness, itching
Eye(s)	Hyperemia, itching, swelling, pain, irritation, foreign body sensation, eye discharge

3. In any of the following cases, please stop using the product, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.
 - (1) If blurriness of the eyes does not improve
 - (2) If symptoms do not improve within approximately 1 week of use in accordance with Dosage and Administration, or you notice any abnormalities.
4. Even if you see improvement of symptoms, consult with your doctor, pharmacist, or registered pharmaceutical distributor if using the product for more than 2 weeks.

Indication

Mitigation of the following eye symptoms:

Drying, foreign body sensation (gritty and prickling sensations), tiredness, blurring, and discomfort while wearing soft or hard contact lenses.

Dosage and administration

Instill 1 drop at a time into the eye, 5 to 6 times a day.

● Please observe the following precautions.


- (1) If administered to a child, the product must be administered under the direction and observation of the child's guardian.
- (2) Do not let the tip of the bottle touch your eyes, eyelids or eyelashes.
- (3) Use only as an eye drop.
- (4) Do not use the product when you are wearing colored contact lenses.

Single-use type

Be sure to dispose of open vials after use, even if there is some solution remaining.

[How to open vials]

Slowly twist the cap to open it, without pulling on it forcefully.




Ingredients and Quantity

Ingredients	Quantity
Purified sodium hyaluronate	0.1%

This product contains the following excipients: aminocaproic acid, edetate sodium hydrate, isotonicizing agent, and pH-adjusting agent.

Precautions for storage and handling

- (1) Store in a cool place away from direct sunlight. To preserve the quality of the product, do not leave it in a hot place such as inside a car or near a heater. If the product is left in a hot place, the bottle may become deformed, resulting in leakage or degradation of the solution. Do not use the product in such a case.
- (2) Keep out of reach of children.
- (3) Do not transfer the solution to another receptacle.
(It may lead to misuse or alter the quality of the product.)
- (4) Do not share the product with another person.
- (5) Placing the product into the back pocket of your trousers or other such places may cause the cap to open. Please exercise caution.
- (6) Do not use the product after the expiry date. Even before the use-by date, please use the product as soon as possible after opening the bottle.



Content Contains 10 vials of 0.3mL