

Disclaimer

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.
- Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

Sante Medical Plus Active®

Precautions



Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

- (1) Patients undergoing medical treatment from a physician
- (2) Patients who have experienced allergic symptoms associated with drugs, etc.
- (3) Persons with the following symptoms:
Severe eye pain
- (4) Persons diagnosed as having the following:
Glaucoma

2. If the following symptoms are observed after using this medicine, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation.

Area affected	Symptom
Skin	Rash/redness, itching
Eyes	Congestion of the eye, itching, swelling, stinging pain

3. In case of the following, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

- (1) When blurred vision does not improve
- (2) When symptoms do not improve even after using the medicine for 5 to 6 days.

Dosage and Administration

Use 1 to 3 drops of this medicine at a time 5 to 6 times a day.

• Please observe the following precautions.

- (1) Excessive use may cause abnormally brightness or eye congestion.
- (2) When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.
- (3) Ensure that the tip of the container does not touch your eyes, eyelids or eyelashes (doing so could cause the solution to become contaminated with eye mucus or bacteria). Do not use when the medicine appears cloudy.
- (4) Do not use this medicine while wearing soft contact lenses.
- (5) Use this medicine only for ophthalmic instillation.



Indication

Blurred vision (increase in eye mucus, etc.), eyestrain, conjunctival congestion, itchy eyes, prevention of eye disease (after swimming, when dust or sweat enters the eyes, etc.), blepharitis (soreness of eyelids), ophthalmia due to light rays such as ultraviolet rays (e.g., snow blindness), discomfort while wearing hard contact lenses

Ingredient and amount

Ingredients	Quantity	Action
Vitamin A (retinol palmitate)	50,000 units /100mL	As well as facilitating secretion of lacrimal fluid (mucin), this product stabilizes tear production and protects the cornea.
Chondroitin sulfate sodium	0.5%	Protects the cornea and moisturizes the eyes by preventing evaporation of tears.
Natural vitamin E (d- α -tocopherol acetate)	0.05%	Improves blood flow in peripheral vessels.
Taurine	0.5%	Activates cell metabolism of eye tissue.
Potassium L-aspartate	0.5%	Increases tissue respiration in the eyes.
Neostigmine methyl sulfate	0.005%	Improves conditions such as eyestrain by improving the focus-adjustment mechanism of the eyes.
Chlorpheniramine maleate	0.03%	Suppresses inflammation and itchiness of the eyes by suppressing histamine activity.
Epsilon-aminocaproic acid	2.0%	Suppresses production of substances that cause inflammation.
Tetrahydrozoline hydrochloride	0.01%	Suppresses congestion of the conjunctiva (the whites of the eyes).

This product contains the following excipients: edetate sodium hydrate, dextran, sodium hyaluronate, hydroxyethyl cellulose, benzalkonium chloride liquid, boric acid, polyoxyethylene hardened castor oil, polysorbate 80, BHT, d-borneol, l-menthol, isotonicizing agent, and pH-adjusting agent.

Precautions for storage and handling

- (1) Store in a cool place away from direct sunlight and keep sealed. To preserve the quality of the product, do not leave it in a hot place such as inside a car or near a heater. If the product is left in a hot place, the vials may become warped causing the solution to leak and degrading its quality. Please do not use the product in such a case.
- (2) Keep out of reach of children.
- (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the medicine.)
- (4) Do not share the product with others.
- (5) Do not use the product if it has exceeded its use-by date. Even before the use-by date, please use the product as soon as possible after opening the bottle.
- (6) Depending on the conditions of storage, the ingredients may crystallize and form white deposits around the inside of the bottle's tip or on the inside of the cap. If this occurs, gently wipe off the deposits with a clean piece of gauze before using the product.

