

## Disclaimer

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.
- Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

# Soft Santear®

## ⚠ Precautions for use



### Consultation

1. The following people should consult with their doctor, pharmacist, or registered pharmaceutical distributor before using this product.
  - (1) People receiving treatment from a doctor
  - (2) People who have experienced an allergic reaction to medicine or other substances
  - (3) People who have the following symptom  
Severe pain in the eyes
  - (4) People who have received the following diagnosis  
Glaucoma
2. If you experience any of the following symptoms after using this product, it may be an adverse reaction. In such a case, please stop using the product immediately, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.

Area affected	Symptom
Skin	Rash/redness, itchiness
Eye(s)	Redness, itchiness, swelling

3. In any of the following cases, please stop using the product, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.
  - (1) If blurriness of the eyes does not improve
  - (2) If symptoms do not improve within approximately 2 weeks of use

## Dosage / Administration

Instill 2 to 3 drops at a time, 5 to 6 times daily.

### ● Please observe the following precautions.


- (1) If administered to a child, the product must be administered under the direction and observation of the child's guardian.
- (2) Do not let the tip of the bottle touch your eyes, eyelids or eyelashes (doing so could cause the solution to become contaminated with eye mucus or bacteria). Do not use the product if it does become contaminated.
- (3) Use only as an eye drop.

## How to open cap

Twist until the cap touches the shoulder of the bottle . . .



A hole is opened in the tip of the bottle.



## Indication

Discomfort while wearing hard or soft contact lenses, supplementing of tear fluid (for dry eyes), tired eyes, bleary eyes (increase in eye mucus, etc.)

## Ingredients and quantity

Ingredients	Quantity
Potassium chloride	0.1%
Sodium chloride	0.4%

This product contains the following excipients: boric acid and pH-adjusting agent.

## Product properties

pH	7.0 to 8.0
Ratio of osmotic pressure	0.90 to 1.20
Properties	Colorless, clear, sterile aqueous eye drop

## Precautions regarding storage and handling

- (1) Do not twist the cap until you are ready to use the product.
- (2) Store in a cool place away from direct sunlight and keep sealed. To preserve the quality of the product, do not leave it in a hot place such as inside a car or near a heater. If the product is left in a hot place, the bottle may become warped causing the solution to leak and degrading its quality. Please do not use the product in such a case.
- (3) Keep out of reach of children.
- (4) Do not transfer the solution to another receptacle.  
(Doing so may lead to misuse of the product or may affect its quality.)
- (5) Do not share the product with another person.
- (6) Do not use the product if it has exceeded its use-by date. Even if it is before the use-by date, please do not use any of the solution remaining in the bottle more than 10 days after opening.
- (7) Depending on the conditions of storage, the ingredients may crystallize and form white deposits around the inside of the bottle's tip or on the inside of the cap. If this occurs, gently wipe off the deposits with a clean piece of gauze before using the product.